HALO®

CANNABIS

Separation of 18 Cannabinoids using HALO® LPH-C18

307



TEST CONDITIONS:

Column: HALO 90 Å LPH-C18, 2.7 µm, 4.6 x 150 mm
Part Number: 92824-716
Mobile Phase A: 5 mM Ammonium Formate, 0.1% Formic Acid
Mobile Phase B: Acetonitrile, 0.1% Formic Acid
Isocratic: 75 %B
Flow Rate: 1.5 mL/min
Pressure: 232 bar
Temperature: 30°C
Detection: PDA, UV: 228 nm
Injection Volume: 3 µL
Sample Solvent: 75/25 MeOH/ Water
Data Rate: 100 Hz
Response Time: 0.025 sec.
Flow Cell: 1 µl
LC System: Shimadzu Nexera X2

PEAK IDENTITIES:

1. Cannabidivarinic acid (CBDVA)
2. Cannabidivarin (CBDV)
3. Cannabidiolic acid (CBDA)
4. Cannabigerolic acid (CBGA)
5. Cannabigerol (CBG)
6. Cannabidiol (CBD)
7. Tetrahydrocannabivarin (THCV)
8. Tetrahydrocannabivarinic acid (THCVA)
9. Cannabinol (CBN)
10. Cannabinolic acid (CBNA)
11. Exo-tetrahydrocannabinol (EXO-THC)
12. delta 9- Tetrahydrocannabinol (D9-THC)
13. delta 8- Tetrahydrocannabinol (D8-THC)
14. Cannabicycol (CBL)
15. Cannabichromene (CBC)
16. Tetrahydrocannabinolic acid A (THCA-A)
17. Cannabichromenic acid (CBCA)
18. Cannabicyclolic acid (CBLA)

A HALO® LPH-C18 column is used to separate a mixture of eighteen cannabinoids, showing fast results and high resolution within critical pairs. Cannabinoids are a class of chemical compounds primarily found in the marijuana plant. Many of these compounds have been found to provide medicinal benefits such as reduction in pain and inflammation.

Peak #	Compound	Transition	CE
1	Carbendazim	192>160.1	-21
2	Dicrotophos	238>112	-22
3	Azamethiphos	324.9>183	-17
4	Pyrimethanil	200.10>107.2	-25
5	Carbofuran	222>123	-22
6	Dodemorph	282.2>116.1	-25
7	Atrazine	216.03>174.1	-17
8	Diuron	232.94>72	-17
9	Iprovalicarb	321.1>119	-30
10	Azoxystrobin	404.04>372.1	-14
11	Fluopram	396.98>208	-25
12	Methoxyfenozide	369.1>149.1	-25
13	Flutolanil	324>242.1	-28
14	Picoxystrobin	368>145.1	-25